

# Saint Mary's Immunization Clinic

Saint Mary's Health and Wellness offers the following immunizations:

CALL 775.770.7100 FOR APPOINTMENTS



**T-dap (Tetanus, Diphtheria and Pertussis)** - This vaccine is required for all students entering seventh grade in both private and public schools. One adult booster is needed for all others 11-64 years of age.

**TD (Tetanus & Diphtheria)**- This vaccine is recommended every ten years to help prevent tetanus for adults 65 and older.

**DTaP (Diphtheria, Tetanus and Pertussis)**- This vaccine is for infants and children under seven years of age. Five doses are needed at two, four, six, 15-18 months and four-six years.

**Hepatitis A**- This vaccine is required for any student attending school in Nevada for the first time. Two doses of this vaccine are needed for lasting protection. These doses should be given at least six months apart. This vaccine is recommended for all travelers, food handlers and daycare providers that pose the risk of spreading the disease to others.

**Hepatitis B**- This vaccine is required for any student attending school in Nevada for the first time. Three doses of this vaccine are needed at zero, one and six months to complete the series.

**HIB (Haemophilus Influenzae Type b)**- This vaccine is for infants and is not routinely recommended after the age of five. Four doses are needed at two, four, six and 12-15 months. The number of doses required depends on what type of vaccine you receive. This is a serious disease caused by a bacteria. It usually strikes children under five years of age.

**HPV (Human Papillomavirus)**- This vaccine is recommended for girls 11-12 years of age and is also recommended for girls and women 13-26 years of age if not already vaccinated. Three doses of this vaccine are needed at zero, two and six months to complete the series. HPV vaccine can prevent 90 percent of genital warts and 70 percent of cervical cancer.

**Influenza**- This vaccine is recommended for all children ages six months through 18 years and all persons 50 years of age and older. Children ages six months through eight years of age receiving the influenza vaccine for the first time should receive two doses at least four weeks apart.

**FluMist**- This is an intranasal influenza vaccine. It is approved for persons two years of age through 49 years of age. Certain criteria must be met in order to receive the intranasal influenza vaccine. Children ages two through eight years of age receiving the vaccine for the first time should receive two doses at least four weeks apart.

**Menactra** This vaccine is recommended for all adolescents 11 through 18 years of age. This vaccine is required for college freshmen living in dormitories.

**MMR (Measles Mumps and Rubella)**- This vaccine is given to all infants and children. Two doses of this vaccine are needed at 12-15 months and four-six years of age. This vaccine is required for all persons attending the University of Nevada, if immunity to these diseases cannot be confirmed.

**Polio (IPV)**- This vaccine is recommended for all children. Four doses of this vaccine are needed at two, four and six-18 months and four-six years of age. People traveling to areas of the world where polio is still common should receive a one time booster dose.

**Pevnar (Pneumococcal Conjugate)**- This vaccine is recommended for all children. Four doses of this vaccine are needed at two, four, six and 12-15 months of age. Pevnar is required for children in daycare.

**Pneumococcal (Pneumonia)**- This vaccine is recommended for all adults 65 years of age and older. The vaccine is also recommended for persons two through 64 years of age who have long-term health problems. New recommendations are also for any adult 19 years of age or older who are cigarette smokers or who have asthma. If you receive a shot before age 65 you will need a booster after age 65 with five years apart.

**Rotavirus**- This vaccine is recommended for all infants. An infant should receive either two or three doses of the vaccine, depending on which brand is used. Rotavirus is a virus that causes severe diarrhea, mostly in babies and young children.

**Twinrix (Hepatitis A & Hepatitis B)**- This vaccine is a combination of the Hepatitis A and Hepatitis B vaccine. It may be given to persons 18 years of age and older. Three doses of this vaccine are needed at zero, one and six months to complete the series.

**Varicella (Chickenpox)**- This vaccine is given to all infants and children. Two doses of this vaccine are needed at 12-15 months and four-five years of age. Anyone 13 years of age and older (who have never had chickenpox or received chickenpox vaccine) should get two doses of the vaccine at least 28 days apart. This vaccine is required for daycare and any student attending school in Nevada for the first time who have not had chickenpox disease.

**Zostavax (Shingles)** This vaccine is recommended for all adults 60 years of age and older. A single dose of this vaccine is needed. Shingles is a painful skin rash, often with blisters. It is also called Herpes Zoster.

*\*A copay is not required for Saint Mary's HealthFirst (HMO/POS) or Health Choice (PPO) members for those vaccines covered under the plans.*



Saint Mary's

A member of CHW

Saint Mary's Immunizations  
343 Elm Street, Suite #203  
Reno, NV 89503  
775.770.3505